




Impact of Grouping Misclassification on Risk Prediction when using the Relative Potency Factors Method to Assess Mixtures Risk

Jeff Swartout, Glenn Rice
U.S. Environmental Protection Agency

Richard C. Hertzberg
Biomathematics Consulting

Society of Toxicology Annual Meeting
San Antonio, TX
March 12, 2018


Disclaimer: The statements and opinions expressed in this presentation are those of the authors and not necessarily those of the U.S. EPA



DISCLAIMER

The authors have no conflicts of interest to report.

The statements and opinions expressed in this presentation are those of the authors and not necessarily those of the U.S. EPA.



ACKNOWLEDGEMENTS

Jane Ellen Simmons, NHEERL

Jason Lambert, NCEA

Marian Olsen, Region 2

David Farrar, NCEA

- Grouping chemicals for mixture risk prediction
 - Extrapolations to environmental exposures
 - In the absence of mixture data
 - Using the index chemical, relative potency factor approach
 - Is it dose addition or response addition (independence)?
- Grouping misclassification effect on mixture risk prediction
 - What can go wrong?
 - How bad is it?
 - Can we fix it?

GROUPING CONCEPTS

- Toxicologic similarity groups for dose addition
 - Similarity of toxic action
 - Common key dose-additive process
 - Not necessarily the full MOA/AOP
- Independent action groups
 - Completely independent MOA (still the same endpoint)
 - No shared key events
 - Dose addition within, response addition across (integrated addition)

INDEX CHEMICAL, RELATIVE POTENCY FACTOR MODEL

IC_RPF = index chemical, RPF dose addition model
Assumes constant relative potency

$$Prob_{mix} = f(\sum_{i=1}^n d_i \cdot RPF_i, \Theta_{IC})$$


$$Prob_{mix} = f(d_{IC} + d_1 \cdot RPF_1 + d_2 \cdot RPF_2 \dots + d_n \cdot RPF_n, \Theta_{IC})$$

IC = index chemical

RPF = relative potency factor

$$RPF_i = EDx_{IC} \div EDx_i$$

EDx = effective dose associated with response x



GROUPING MISCLASSIFICATION: Where can we go wrong?

- Getting the RPF wrong
- Assigning a compound to the wrong dose-additive group
- Failing to allow for differently-shaped dose-response curves
- Mistaking dose additivity for response additivity
- Failing to account for different control values
- Failing to account for partial agonists
- Improper application of effect summation
- Low-dose extrapolation


GROUPING MISCLASSIFICATION I: Getting the RPF wrong

➤ Getting the RPF wrong

- Error depends on the magnitude of the RPF error, the mass-fraction contribution of the chemical to the mixture, and the group IC dose-response function
- Assigning a compound to the wrong dose-additive group
- Failing to allow for differently-shaped dose-response curves
- Mistaking dose additivity for response additivity
- Failing to account for different control values
- Failing to account for partial agonists
- Improper application of effect summation
- Low-dose extrapolation

GROUPING MISCLASSIFICATION I: Getting the RPF wrong

Chemical components of mixture
Weibull power = 1




GROUPING MISCLASSIFICATION I: Getting the RPF wrong

2-chemical mixture of equal doses

[Chem2]:[Chem1] = 1

Weibull power = 1




GROUPING MISCLASSIFICATION I: Getting the RPF wrong

2-chemical mixture of equi-toxic doses


[Chem2]:[Chem1] = 100

Weibull power = 1



GROUPING MISCLASSIFICATION I: Getting the RPF wrong

Chemical components of mixture
Weibull power = 2




GROUPING MISCLASSIFICATION I: Getting the RPF wrong

2-chemical mixture of equi-toxic doses

[Chem2]:[Chem1] = 100

Weibull power = 1




GROUPING MISCLASSIFICATION I: Getting the RPF wrong

2-chemical mixture of equal doses

[Chem2]:[Chem1] = 1

Weibull power = 2




GROUPING MISCLASSIFICATION I: Getting the RPF wrong

10-chemical mixture of equi-toxic doses

[Chem2]:[Chem1] = 10

Weibull power = 2



GROUPING MISCLASSIFICATION II: Wrong Group Assignment

- Getting the RPF wrong
- **Assigning a compound to the wrong dose-additive group**
 - Applies to integrated addition of independent action groups
 - Error depends on the relative potency of the chemical, the mass-fraction contribution of the chemical to the group, and the group IC dose-response function
- Failing to allow for differently-shaped dose-response curves
- Mistaking dose additivity for response additivity
- Failing to account for different control values
- Failing to account for partial agonists
- Improper application of effect summation
- Low-dose extrapolation




GROUPING MISCLASSIFICATION III: Dose-Response Curve Shape

- Getting the RPF wrong
- Assigning a compound to the wrong dose-additive group
- **Failing to allow for differently-shaped dose-response curves**
- Mistaking dose additivity for response additivity
- Failing to account for different control values
- Failing to account for partial agonists
- Improper application of effect summation
- Low-dose extrapolation

DOSE-RESPONSE CURVE SHAPE


- Same shape required (mathematically) for constant relative potency at all doses and full applicability of IC_RPF model
- Different shape does not mean dose addition does not apply (as previously interpreted by many)
 - High-dose non-proportionalities in the toxicokinetics, but additivity at a common molecular initiating event (MIE)
- Failing to allow for differently-shaped dose-response curves can result in potentially significant error when using the IC_RPF method
 - Assuming same shape when they are different
 - Assuming different shapes when they are the same
 - Mistaking dose additivity for response additivity

- Simple similar action at the MIE
- Proportionality in toxicokinetics & toxicodynamics
 - RPF-scaled external doses are additive (constant relative potency)




*WP = Weibull power

DOSE ADDITION FOR CHEMICALS WITH SIMILAR DOSE-RESPONSE SHAPES




- Mixture for any combination of fixed-proportion doses will have same shape

- Simple similar action at the MIE
- Non-proportionalities in toxicokinetics
 - RPF-scaled external doses are **not** additive (non-constant relative potency)
 - Predicted mixture risk dependent on choice of IC (Chen et al., 2003)
 - Addition of internal component doses




*WP = Weibull power

DOSE ADDITION FOR CHEMICALS WITH DISSIMILAR DOSE-RESPONSE SHAPES



- Mixture response prediction is dependent on the choice of IC


DOSE ADDITION FOR CHEMICALS WITH DISSIMILAR DOSE-RESPONSE SHAPES



- Mixture response prediction is dependent on the choice of IC

VARIABLE RELATIVE POTENCY: IC DEPENDENCE ERROR

Two component mixture: component dose-response shape parameters are 2-fold different



GROUPING MISCLASSIFICATION IV: Dose Additivity vs. Response Additivity

- Getting the RPF wrong
- Assigning a compound to the wrong dose-additive group
- Failing to allow for differently-shaped dose-response curves
- **Mistaking dose additivity for response additivity**
 - Failing to account for different control values
 - Failing to account for partial agonists
 - Improper application of effect summation
 - Low-dose extrapolation

DOSE ADDITION INTERPRETED AS RESPONSE ADDITION

- The case of differently-shaped component dose-response curves (DRCs)
 - If one assumes Bliss “dilution” concept applies to both toxicokinetics and toxicodynamics
- The case of different MIEs
 - Could be assumed to be evidence for total independence of toxicodynamic pathways
- Result is likely to be under-predictive
 - Except for thresholds


DOSE ADDITION INTERPRETED AS RESPONSE ADDITION

- The case of differently-shaped component dose-response curves (DRCs)
 - If one assumes Bliss “dilution” concept applies to both toxicokinetics and toxicodynamics
- The case of different MIEs
 - Could be assumed to be evidence for total independence of toxicodynamic pathways
- Result is likely to be under-predictive

DIFFERENT MIEs

- Often interpreted as definitive for independence and use of response addition
- Dose addition, however, can apply for downstream concentrations of common or similarly-acting products in the toxicodynamic pathways (shared dose-additive event).
 - Hormone-driven outcomes (androgen insufficiency; Rider et al., 2010)
 - Reactive oxygen species toxicity (PaH's ?)
 - Ion channel blockers (Norberg & Wahlstrom, 1988 ?)

DOSE ADDITION FOR INITIALLY INDEPENDENT PATHWAYS



Independent MIE suggests response addition may apply

Pathway convergence at “dose”-additive process may allow for dose addition

Modification of Lambert and Lipscomb (2007)

GROUPING MISCLASSIFICATION V: Continuous Measures

- Getting the RPF wrong
- Assigning a compound to the wrong dose-additive group
- Failing to allow for differently-shaped dose-response curves
- Mistaking dose additivity for response additivity
- **Failing to account for different control values**
- **Failing to account for partial agonists**
- **Improper application of effect summation**
- Low-dose extrapolation

GROUPING MISCLASSIFICATION VI: Low-dose Extrapolation

- Getting the RPF wrong
- Assigning a compound to the wrong dose-additive group
- Failing to allow for differently-shaped dose-response curves
- Mistaking dose additivity for response additivity
- Failing to account for different control values
- Failing to account for partial agonists
- Improper application of effect summation
- **Low-dose extrapolation**
 - All mixture risk models assume some form of continuation of component dose-response curve shape below the tested range, but how can we know?
 - Need a way to inform low-dose mixture shape prediction *experimentally*

SUMMARY & CONCLUSIONS

- Grouping for dose additivity starts with similarity of toxic action
 - Key dose-additive event is required, but does not have to be the MIE
- Dose addition can still apply with dissimilar dose-response curve shapes
 - IC_RPF model is impacted adversely
- Dose addition can still apply with independent MIEs
 - Downstream toxicodynamic pathway convergence
- Grouping errors can be substantial but could be “diluted” out with large number of chemicals
- Low-dose extrapolation is a major issue
 - Need a way to predict dose-response shape below the tested range (other than mathematical speculation)
 - How about some of that 21st Century Toxicology?



REFERENCES

- Chen, JJ; Chen, Y-J; Teuschler, LK; Rice, G; Hamernik, K; Protzel, A; Kodell, RL. (2003). Cumulative risk assessment for quantitatative response date. Environmetrics 14: 339-353.
- Gennings, C; Carter, WH, Jr.; Carney, EW; Charles, GD; Gollapudi, BB; Carchman, RA. (2004). A novel flexible approach for evaluating fixed ratio mixtures of full and partial agonists. Toxicol Sci 80: 134-150. <http://www.ncbi.nlm.nih.gov/pubmed/15084752>
- Lambert, J; Lipscomb J. (2007). Mode of action as a determining factor in additivity models for chemical mixture risk assessment. Toxicol Appl Pharmacol 14: 340-347
- National Research Council (NRC) (2007). Phthalates and Cumulative Risk Assessment The Task Ahead. Committee on the Health Risks of Phthalates, National Research Council. National Academies Press. <http://www.nap.edu/catalog/12528.html>
- Rider, CV; Furr, JR; Wilson, VS; Gray, LE, Jr. (2010). Cumulative effects of in utero administration of mixtures of reproductive toxicants that disrupt common target tissues via diverse mechanisms of toxicity. Int J Androl 33: 443-462. <http://www.ncbi.nlm.nih.gov/pubmed/20487044>
- U.S. EPA (Environmental Protection Agency). (2000). Supplementary guidance for conducting health risk assessment of chemical mixtures (pp. 209). (EPA/630/R-00/002). Washington, DC: Risk Assessment Forum. http://ofmpub.epa.gov/eims/eimscomm.getfile?p_download_id=4486
- U.S. EPA (Environmental Protection Agency). (2016) Pesticide Cumulative Risk Assessment: Framework for Screening Analysis Purpose. Office of Pesticide Programs, Office of Chemical Safety and Pollution Prevention